


GDAS atmospheric models in astroparticle shower simulations




J. Grisales-Casadiegos¹, C. Sarmiento-Cano², Luis A. Núñez¹, for the LAGO Collaboration
¹Universidad Industrial de Santander, Colombia, ²Instituto en Tecnologías de Detección y Astropartículas, Argentina


How the atmosphere influences the background flux of secondary Cosmic Rays?




The LAGO collaboration exploits its array of detectors to examine the influence of space weather on the cosmic ray flux at the Earth surface.



We observed some differences between the GDAS and the CORSIKA default models (tropical and sub-tropical summer) in the first 30 km.





RESULTS




The secondary particle flux at ground level are composed by different particles that dominates at certain energies. We observed the known flux distribution with the new atmospheric profiles.

HOW WE DO THAT?

Creating monthly atmospheric profiles



We observed some differences between April and November, the latter being the closest to the sub-tropical summer model.



SUMMARY

- We have **devised** a methodology that enables one to obtain a month-by-month averaged atmospheric profiles for any geographic location.
- These results **suggest** the importance of continuing to study these effects in greater detail.
- Through this work we **update** the sequence of simulations for the LAGO collaboration, to study the modulation of the solar wind on the flux of secondaries detected.

Scan for more information:



Questions or suggestions: jennifer.grisales@saber.uis.edu.co

Complete list of authors:



GDAS atmospheric models in astroparticle shower simulations





Universidad Industrial de Santander

J. Grisales-Casadiegos¹, C. Sarmiento-Cano², Luis A. Núñez¹, for the LAGO Collaboration
¹Universidad Industrial de Santander, Colombia, ²Instituto en Tecnologías de Detección y Astropartículas, Argentina

How the atmosphere influences the background flux of secondary Cosmic Rays?

The LAGO collaboration exploits its array of detectors to examine the influence of space weather on the cosmic ray flux at the Earth surface.



Here, we used GDAS (Global Data Assimilation System) to study the impact of climate variability in the background secondary flux.




GDAS atmospheric models in astroparticle shower simulations




J. Grisales-Casadiegos¹, C. Sarmiento-Cano², Luis A. Núñez¹, for the LAGO Collaboration
¹Universidad Industrial de Santander, Colombia, ²Instituto en Tecnologías de Detección y Astropartículas, Argentina


How the atmosphere influences the background flux of secondary Cosmic Rays?




The LAGO collaboration exploits its array of detectors to examine the influence of space weather on the cosmic ray flux at the Earth surface.



We observed some differences between the GDAS and the CORSIKA default models (tropical and sub-tropical summer) in the first 30 km.





RESULTS




The secondary particle flux at ground level are composed by different particles that dominates at certain energies. We observed the known flux distribution with the new atmospheric profiles.

HOW WE DO THAT?

Creating monthly atmospheric profiles



We observed some differences between April and November, the latter being the closest to the sub-tropical summer model.



SUMMARY

- We have **devised** a methodology that enables one to obtain a month-by-month averaged atmospheric profiles for any geographic location.
- These results **suggest** the importance of continuing to study these effects in greater detail.
- Through this work we **update** the sequence of simulations for the LAGO collaboration, to study the modulation of the solar wind on the flux of secondaries detected.

Scan for more information:



Questions or suggestions:
jennifer.grisales@saber.uis.edu.co

Complete list of authors:



GDAS atmospheric models in astroparticle shower simulations





Universidad Industrial de Santander

J. Grisales-Casadiegos¹, C. Sarmiento-Cano², Luis A. Núñez¹, for the LAGO Collaboration

¹Universidad Industrial de Santander, Colombia, ²Instituto en Tecnologías de Detección y Astropartículas, Argentina

HOW WE DO THAT?

Creating monthly atmospheric profiles




GDAS atmospheric models in astroparticle shower simulations




J. Grisales-Casadiegos¹, C. Sarmiento-Cano², Luis A. Núñez¹, for the LAGO Collaboration
¹Universidad Industrial de Santander, Colombia, ²Instituto en Tecnologías de Detección y Astropartículas, Argentina


How the atmosphere influences the background flux of secondary Cosmic Rays?




The LAGO collaboration exploits its array of detectors to examine the influence of space weather on the cosmic ray flux at the Earth surface.



We observed some differences between the GDAS and the CORSIKA default models (tropical and sub-tropical summer) in the first 30 km.





RESULTS




The secondary particle flux at ground level are composed by different particles that dominates at certain energies. We observed the known flux distribution with the new atmospheric profiles.

HOW WE DO THAT?

Creating monthly atmospheric profiles



We observed some differences between April and November, the latter being the closest to the sub-tropical summer model.



SUMMARY

- We have **devised** a methodology that enables one to obtain a month-by-month averaged atmospheric profiles for any geographic location.
- These results **suggest** the importance of continuing to study these effects in greater detail.
- Through this work we **update** the sequence of simulations for the LAGO collaboration, to study the modulation of the solar wind on the flux of secondaries detected.

Scan for more information:



Questions or suggestions:
 jennifer.grisales@saber.uis.edu.co

Complete list of authors:



GDAS atmospheric models in astroparticle shower simulations




Universidad Industrial de Santander


J. Grisales-Casadiegos¹, C. Sarmiento-Cano², Luis A. Núñez¹, for the LAGO Collaboration

¹Universidad Industrial de Santander, Colombia, ²Instituto en Tecnologías de Detección y Astropartículas, Argentina

We observed some differences between the GDAS and the CORSIKA default models (tropical and sub-tropical summer) in the first 30 km.



RESULTS



The secondary particle flux at ground level are composed by different particles that dominates at certain energies. We observed the known flux distribution with the new atmospheric profiles.




GDAS atmospheric models in astroparticle shower simulations




J. Grisales-Casadiegos¹, C. Sarmiento-Cano², Luis A. Núñez¹, for the LAGO Collaboration
¹Universidad Industrial de Santander, Colombia, ²Instituto en Tecnologías de Detección y Astropartículas, Argentina


How the atmosphere influences the background flux of secondary Cosmic Rays?




The LAGO collaboration exploits its array of detectors to examine the influence of space weather on the cosmic ray flux at the Earth surface.



We observed some differences between the GDAS and the CORSIKA default models (tropical and sub-tropical summer) in the first 30 km.





RESULTS




The secondary particle flux at ground level are composed by different particles that dominates at certain energies. We observed the known flux distribution with the new atmospheric profiles.

HOW WE DO THAT?

Creating monthly atmospheric profiles



We observed some differences between April and November, the latter being the closest to the sub-tropical summer model.



SUMMARY

- We have **devised** a methodology that enables one to obtain a month-by-month averaged atmospheric profiles for any geographic location.
- These results **suggest** the importance of continuing to study these effects in greater detail.
- Through this work we **update** the sequence of simulations for the LAGO collaboration, to study the modulation of the solar wind on the flux of secondaries detected.

Scan for more information:



Questions or suggestions:
 jennifer.grisales@saber.uis.edu.co

Complete list of authors:



GDAS atmospheric models in astroparticle shower simulations




Universidad Industrial de Santander

J. Grisales-Casadiegos¹, C. Sarmiento-Cano², Luis A. Núñez¹, for the LAGO Collaboration

¹Universidad Industrial de Santander, Colombia, ²Instituto en Tecnologías de Detección y Astropartículas, Argentina

We observed some differences between April and November, the latter being the closest to the sub-tropical summer model.




GDAS atmospheric models in astroparticle shower simulations




J. Grisales-Casadiegos¹, C. Sarmiento-Cano², Luis A. Núñez¹, for the LAGO Collaboration
¹Universidad Industrial de Santander, Colombia, ²Instituto en Tecnologías de Detección y Astropartículas, Argentina


How the atmosphere influences the background flux of secondary Cosmic Rays?




The LAGO collaboration exploits its array of detectors to examine the influence of space weather on the cosmic ray flux at the Earth surface.



We observed some differences between the GDAS and the CORSIKA default models (tropical and sub-tropical summer) in the first 30 km.





RESULTS




The secondary particle flux at ground level are composed by different particles that dominates at certain energies. We observed the known flux distribution with the new atmospheric profiles.

HOW WE DO THAT?

Creating monthly atmospheric profiles



We observed some differences between April and November, the latter being the closest to the sub-tropical summer model.



SUMMARY

- We have **devised** a methodology that enables one to obtain a month-by-month averaged atmospheric profiles for any geographic location.
- These results **suggest** the importance of continuing to study these effects in greater detail.
- Through this work we **update** the sequence of simulations for the LAGO collaboration, to study the modulation of the solar wind on the flux of secondaries detected.

Scan for more information:



Questions or suggestions: jennifer.grisales@saber.uis.edu.co

Complete list of authors:



GDAS atmospheric models in astroparticle shower simulations

J. Grisales-Casadiegos¹, C. Sarmiento-Cano², Luis A. Núñez¹, for the LAGO Collaboration

¹Universidad Industrial de Santander, Colombia, ²Instituto en Tecnologías de Detección y Astropartículas, Argentina




Universidad
Industrial de
Santander

SUMMARY

- We have **devised** a methodology that enables one to obtain a month-by-month averaged atmospheric profiles for any geographic location.
- These** results **suggest** the importance of continuing to study these effects in greater detail.
- Through** this work we **update** the sequence of simulations for the LAGO collaboration, to study the modulation of the solar wind on the flux of secondaries detected.

Scan for more
information:



Complete list
of authors:



Questions or suggestions:
jennifer.grisales@saber.uis.edu.co

ONLINE ICRC 2021

